

AN UNUSUAL PROOF THAT THE REALS ARE UNCOUNTABLE

ELIAHU LEVY

ABSTRACT. This somewhat unusual proof for the fact that the reals are uncountable, which is adapted from one of Bourbaki's proofs in "Fonctions d'une variable réelle", may be of some interest.

Claim . *The set $\mathbb{R}$ of the real numbers is uncountable.*

Proof . (adapted from a proof in [B]). Suppose $\mathbb{R}$ was countable. Then there is a function $a(x) : \mathbb{R} \rightarrow \mathbb{R}$ such that:

1. $a(x) > 0$ for all x (strictly positive),
 2. the sum of the $a(x)$ on any finite set is ≤ 1 .
- (just take $a(x) = 2^{-n}$ if x is the n 'th element in the counting).

Now, define for any subset S of $\mathbb{R}$,

$m(S) :=$ the supremum of the sums of $a(x)$ on finite subsets of S .

Then surely $0 \leq m(S) \leq 1$ for any S . (In fact, $m(S)$ will be used only for S an open ray.)

For $x \in \mathbb{R}$, compare $m] - \infty, x[$ (open ray) with x . One may define:

$$c := \sup \left\{ x \mid m] - \infty, x[> x \right\},$$

and we shall reach a contradiction.

Indeed, since $a(c) > 0$, there is a y such that $y > c - a(c)$ and $m] - \infty, y[> y$, thus $y \leq$ than the supremum c . Now, since $y \leq c$, $] - \infty, y[$ does not contain c . But $] - \infty, y + a(c)[$ contains c because $y > c - a(c)$. So by the definition of $m(S)$,

$$m] - \infty, y + a(c)[\geq m] - \infty, y[+ a(c) > y + a(c),$$

yet $y + a(c)$ exceeds the supremum c , a contradiction. □

REFERENCES

[B] N. Bourbaki, *Fonctions d'une variable réelle*. Diffusion C. C. L. S., Paris.

DEPARTMENT OF MATHEMATICS, TECHNION – ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000, ISRAEL
E-mail address: eliahu@techunix.technion.ac.il